

On the connection between financial processes with stochastic volatility and nonextensive statistical mechanics

S. M. Duarte Queirós¹ and C. Tsallis^{1,2 *}

¹*Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud 150,
22290-180, Rio de Janeiro-RJ, Brazil*

²*Santa Fe Institute, 1399 Hyde Park Road,
Santa Fe, New Mexico 87501, USA*

(February 2, 2008)

The *GARCH* algorithm is the most renowned generalisation of Engle's original proposal for modelling *returns*, the *ARCH* process. Both cases are characterised by presenting a time dependent and correlated variance or *volatility*. Besides a memory parameter, b , (present in *ARCH*) and an independent and identically distributed noise, ω , *GARCH* involves another parameter, c , such that, for $c = 0$, the standard *ARCH* process is reproduced. In this manuscript we use a generalised noise following a distribution characterised by an index q_n , such that $q_n = 1$ recovers the Gaussian distribution. Matching low statistical moments of *GARCH* distribution for returns with a q -Gaussian distribution obtained through maximising the entropy $S_q = \frac{1 - \sum_i p_i^q}{q-1}$, basis of nonextensive statistical mechanics, we obtain a sole analytical connection between q and (b, c, q_n) which turns out to be remarkably good when compared with computational simulations. With this result we also derive an analytical approximation for the stationary distribution for the (squared) volatility. Using a generalised Kullback-Leibler relative entropy form based on S_q , we also analyse the degree of dependence between successive returns, z_t and z_{t+1} , of *GARCH*(1, 1) processes. This degree of dependence is quantified by an entropic index, q^{op} . Our analysis points the existence of a unique relation between the three entropic indexes q^{op} , q and q_n of the problem, independent of the value of (b, c) .

I. INTRODUCTION

The study of time series plays an important role in science due to their ubiquity in both natural and artificial systems. They can be found, e.g., in geoseismic (earthquakes), meteorological (El Niño), physiological (electroencephalographic profiles) or financial phenomena [1–5]. They comprise a sequentially ordered set of random variables, correlated or not, following a certain probability function. For the uncorrelated case, the most perceptive is to consider the time series as a succession of values that are associated to the same probability distribution, like it occurs for the ordinary random walk, where probability of a certain jump value is constant in time. This kind of process is defined as *homoskedastic*. However, there are phenomena for which the probability distribution associated to the stochastic variable at some time step t depends explicitly on t and these are named *heteroskedastic*. A simple way to obtain a heteroskedastic process is to consider the same probability functional for all times, but with a varying second-order moment (or width). In this sort of stochasticity we can include financial time series, namely return time series, where second-order moment time dependence is a fea-

ture more than well-known [6]. Aiming to mimic this type of systems, Engle introduced in 1982 the autoregressive conditional heteroskedasticity (*ARCH*(s)) (s will be defined later on) process [7], which is considered a landmark in finance, comparable to the Black-Scholes equation [8], because of its wide use [9,10]. Albeit its extraordinary importance, Engle's model presents, in some situations, implementation difficulties and above that, is not able to fully satisfactorily reproduce the empirically observed power-law like correlation decay of volatility. These failures lead T. Bollerslev to generalise it defining the *GARCH*(s, r) [11] (G stands for *generalised*). Due to its financial cradle these models are not well-known in physics, nevertheless they can be very useful to illustrate many traditional physical problems (see, e.g., [12]). In this manuscript we give sequence to the ansatz presented by us in a previous work [13] for the *ARCH*(1) process, generalising it to the *GARCH*(1, 1). Moreover, we detail the physical justification for why these models (which present time dependent variance), accommodate well within the current nonextensive statistical mechanical theory. We also present an analytical form for the distribution for the (squared) volatility and analyse the degree of dependence between successive *returns*. The

*E-mail addresses: sdqueiro@cbpf.br, tsallis@santafe.edu

manuscript is organised as follows: In Sec. II we introduce the $GARCH(s,r)$ process and some of its properties; In Sec. III A we present our connection between $GARCH(1,1)$ and nonextensive entropy. In addition, along the lines of superstatistics [15], we derive the distribution for the (squared) volatility in Sec. III B. In Sec. IV, applying a generalised Kullback-Leibler relative entropy we analyse the degree of dependence between z_t and z_{t+1} elements of a $GARCH(1,1)$ time series and its relation with non-Gaussianity. Some final comments are made in Sec. V.

II. THE GARCH MODEL

As settled by Engle [7], we will define an autoregressive conditional heteroskedastic time series as a discrete time stochastic process, z_t ,

$$z_t = \sigma_t \omega_t, \quad (1)$$

where ω_t is an independent and identically distributed random variable with null mean and unitary variance, i.e., $\langle \omega_t \rangle = 0$ and $\langle \omega_t^2 \rangle = 1$. The quantity σ_t , named *volatility* is time varying, positive defined and dependent of the past values of the *return* z_t . According to its definition, the process presents mean zero, is uncorrelated ($\langle z_t z_{t'} \rangle \sim \delta_{tt'}$) and has a conditional variance, σ_t^2 , that evolves in time.

In his original paper Engle [7] suggests a possible expression for σ_t^2 defining it as a linear function of past squared values of z_t known as *ARCH(s) linear process*,

$$\sigma_t^2 = a + \sum_{i=1}^s b_i z_{t-i}^2, \quad (a, b_i \geq 0). \quad (2)$$

With the aim of solving the weak points of the *ARCH* process that were referred in Sec. I, it was introduced the linear $GARCH(s,r)$, which presents a more flexible structure for the functional form of σ_t^2 decreasing the direct influence of z on σ_t^2 (for details see [11]):

$$\sigma_t^2 = a + \sum_{i=1}^s b_i z_{t-i}^2 + \sum_{i=1}^r c_i \sigma_{t-i}^2, \quad (a, b_i, c_i \geq 0), \quad (3)$$

Like its predecessor, the $GARCH(s,r)$ model also captures the recognised tendency for *volatility clustering* (evident in financial time series) and is very similar to intermittent fluctuations in turbulent flows [14]: large (small) values of z_t are usually followed by large (small) values. However, due to the *arbitrary sign* of ω_t , it can be verified that, although $\langle z_t z_{t'} \rangle \sim \delta_{tt'}$, $\langle |z_t| |z_{t'}| \rangle$ is *not* proportional to $\delta_{tt'}$. For $c_i = 0 (\forall i)$, $GARCH(s,r)$ straightforwardly reduces to the linear *ARCH(s)*. See Figs. 1, 2 and 3 for typical realisations.




FIG. 1. Typical $GARCH(1,1)$ time series obtained for a Gaussian noise, $q_n = 1$. In (a) we present time series which correspond to pure Gaussian ($\square$) and *ARCH(1)* ($\circ$). In (b) the introduction of parameter c ($\diamond$) increases the probability for larger values of $|z_t|$, thus increasing tails in $P(z)$.




FIG. 2. Time dependence of volatility σ for the $GARCH$ processes presented in Fig. 1. Here is clearly visible the difference between $c = 0$ (*ARCH*) and $c \neq 0$ (*GARCH*). For the same values of b we are able to obtain greater values of σ , thus leading to fatter tails in $P(z)$.




FIG. 3. In (a) correlation between returns $z(t+T)$ and $z(t)$ vs. time lag T . All values, except for $T = 0$, are at noise level. In (b) correlation between absolute values of returns which present a decay of time lag T . The run analysed was the same presented in Fig. 1

Let us focus on the simplest and most used of the *GARCH* processes, namely the *GARCH*(1,1) one. In this case,

$$\sigma_t^2 = a + b z_{t-1}^2 + c \sigma_{t-1}^2, \quad (4)$$

so that the process is completely defined when a , b , c , and the noise nature, $P_n(\omega_t)$, are specified.

Combining Eqs. (1) and (4) we obtain the n^{th} order moment for the stationary $P(z)$ distribution, particularly the second,

$$\bar{\sigma}^2 \equiv \langle z_t^2 \rangle = \langle \sigma_t^2 \rangle = \frac{a}{1 - b - c}, \quad (b + c < 1), \quad (5)$$

and the fourth moment,

$$\langle z_t^4 \rangle = a^2 \langle \omega_t^4 \rangle \frac{1 + b + c}{(1 - b - c)(1 - 2bc - c^2 - b^2 \langle \omega_t^4 \rangle)}. \quad (6)$$

Let us now assume, for simplicity and without lack of generality, a *GARCH*(1,1) process that generates time series with unitary variance, i.e., $\bar{\sigma}^2 = 1$, which imposes $a = 1 - b - c$. Now, for this process, the fourth moment is numerically equal to the kurtosis $k_x \equiv \frac{\langle x^4 \rangle}{\langle x^2 \rangle^2}$, and thus we get,

$$\langle z_t^4 \rangle = k_z = k_\omega \left(1 + b^2 \frac{k_\omega - 1}{1 - c^2 - 2bc - b^2 k_\omega} \right), \quad (7)$$

where $c^2 + 2bc + b^2 k_\omega < 1$. It is clear from Eq. (7) that *GARCH*(1,1) generates distributions $P(z)$ with fatter

tails than those of the noise ω_t , and fatter also than the ones obtained with *ARCH*(1) with the same b [16]. It is also possible to see that parameter c is only useful for $b \neq 0$, otherwise *GARCH*(1,1) process reduces to constructing stationary probability distributions with a kurtosis k_z equal to k_ω .

III. THE ANSATZ CONNECTING GARCH AND NONEXTENSIVE STATISTICAL MECHANICS

A. Stationary distribution for returns

We shall now establish a connection — physically motivated in the next subsection — between the present stochastic process and the current nonextensive statistical mechanical framework, based on the entropic form [17],

$$S_q = \frac{1 - \int_{-\infty}^{+\infty} [p(z)]^q dz}{q - 1}, \quad (q \in \mathbb{R}). \quad (8)$$

This entropy is currently referred to as *nonextensive* because it is so for *independent* subsystems. It can however be *extensive* in the presence of scale-invariant correlations [18–21]. The associated statistics has been successfully applied to phenomena presenting some kind of scale-invariant geometry, like in low-dimensional dissipative and conservative maps [22], anomalous (correlated) diffusion [23], turbulent flows [24], Langevin dynamics with fluctuating temperature [15,25,26], long-range many-body classical Hamiltonians [27], among many others [28]. Entropy (8) constitutes a generalisation of the Boltzmann-Gibbs (BG) one, namely $S_{BG} = - \int_{-\infty}^{+\infty} p(z) \ln p(z) dz$. Indeed, this celebrated expression is recovered as the $q \rightarrow 1$ limit of entropy (8).

By applying the standard variational principle on entropy (8) with the constraints $\int_{-\infty}^{+\infty} p(z) dz = 1$ and $\int_{-\infty}^{+\infty} z^2 [p(z)]^q dz / \int_{-\infty}^{+\infty} [p(z)]^q dz = \bar{\sigma}_q^2$ [29–31] ($\bar{\sigma}_q^2$ is defined as the *generalised second-order moment*) we obtain

$$p(z) = \frac{\mathcal{A}}{[1 + \mathcal{B} (q - 1) z^2]^{\frac{1}{q-1}}}, \quad \left(q < \frac{5}{3} \right), \quad (9)$$

where

$$\mathcal{B} \equiv \frac{1}{\bar{\sigma}^2 (5 - 3q)}, \quad (10)$$

$$\bar{\sigma}^2 \equiv \int_{-\infty}^{+\infty} p(z) dz, \quad (11)$$

and

$$\mathcal{A} = \frac{\Gamma\left[\frac{1}{q-1}\right]}{\sqrt{\pi} \Gamma\left[\frac{3-q}{2q-2}\right]} \sqrt{(q-1) \mathcal{B}}. \quad (12)$$

Standard and generalised second-order moments, $\bar{\sigma}^2$ and $\bar{\sigma}_q^2$, are related by $\bar{\sigma}^2 = \bar{\sigma}_q^2 \frac{3-q}{5-3q}$. In the limit $q \rightarrow 1$, distribution (9) becomes a Gaussian. If $q = \frac{3+m}{1+m}$ ($m = 1, 2, 3, \dots$), distribution (9) recovers the Student's t -distribution with m degrees of freedom; if $q = \frac{n-4}{n-2}$ ($n = 3, 4, 5, \dots$), it recovers the so-called r -distribution with n degrees of freedom [32].

Defining the q -exponential function as,

$$\exp_q(x) \equiv [1 + (1-q)x]^{\frac{1}{1-q}} \quad (\exp_1(x) = e^x), \quad (13)$$

we can rewrite the above distribution as follows,

$$p(z) = \mathcal{A} e_q^{-Bz^2}, \quad (14)$$

from now on referred to as q -Gaussian. The fourth moment of $p(z)$ is,

$$\langle z^4 \rangle = 3 (\bar{\sigma}^2)^2 \frac{3q-5}{5q-7}. \quad (15)$$

Let us now propose the ansatz $p(z) \simeq P(z)$ with $\bar{\sigma}^2 = 1$. Specifically, we will impose the matching of Eqs. (7) and (15). We assume that the noise ω_t follows the generalised distribution

$$P_n(\omega) = \frac{\mathcal{A}_{q_n}}{\left[1 + \frac{q_n-1}{5-3q_n} \omega^2\right]^{\frac{1}{q_n-1}}}, \quad \left(q_n < \frac{5}{3}\right), \quad (16)$$

defined by the index q_n ; its variance equals unity, and $\mathcal{A}_{q_n}$ is uniquely determined through normalization. We are then able to establish a relation between parameters b , c and indices q_n and q :

$$b = \frac{\sqrt{(q-q_n)[(5-3q_n)(2-q)-c^2(5-3q)(2-q_n)]}}{(5-3q_n)(2-q)} - \frac{c(q-q_n)}{(5-3q_n)(2-q)}. \quad (17)$$

For $b = c = 0$ we have $q = q_n$. Naturally, the connection indicated in Eq. (17) and illustrated in Fig. 4 for typical values of q_n reduces, for $c = 0$, to the one obtained [13] for the ARCH(1) model, namely $b = \sqrt{(q-q_n)/[(5-3q_n)(2-q)]}$, hence $q = [q_n + 2b^2(5-3q_n)]/[1 + b^2(5-3q_n)]$.

To verify the above ansatz we generated, for typical values of (b, c, q_n) and using an algorithm based on Eqs. (1) and (4), a set of GARCH time series. Then we computed the corresponding probability density functions and compared them with the histograms (with any adequately chosen interval δ) associated with the q -Gaussian distribution with q satisfying the ansatz. We compared then the numerical probability density functions (PDFs) with $H(z)$

$$H(z) = \int_{z+\delta/2}^{z+\delta/2} p(x) dx = \frac{\Gamma[\frac{1}{q-1}]}{2\Gamma[\frac{1}{q-1}-\frac{1}{2}]} \sqrt{\frac{1-q}{\pi(3q-5)}} \times \left\{ (\delta-2z) {}_2F_1\left(\frac{1}{2}, \frac{1}{q-1}; \frac{3}{2}; \frac{(q-1)(\delta-2z)^2}{4(3q-5)}\right) + (\delta+2z) {}_2F_1\left(\frac{1}{2}, \frac{1}{q-1}; \frac{3}{2}; \frac{(q-1)(\delta+2z)^2}{4(3q-5)}\right) \right\}, \quad (18)$$

where ${}_2F_1$ is the hypergeometric function.




FIG. 4. Diagram (q, b, c) for $q_n = 1$ in a) and $q_n = 1.2$ in b). In b) we can see that the greatest allowed value of b ($c = 0$) is $b = \frac{1}{\sqrt{4.2}} \simeq 0.488$.

As can be seen in Figs. 5 and 6, the accordance between stationary PDFs and the PDFs obtained by using, in Eq. (9), the value of q satisfying Eq. (17), is quite satisfactory. In the captions of Figs. 5 and 6, we present the values of the χ^2 error function,

$$\chi^2 \equiv \frac{1}{N} \sum_{i=1}^N [P(z) - H(z)]^2.$$

Another way to evaluate the slight discrepancy between $P(z)$ and $p(z)$ (or $H(z)$) is to compute the percentual error in the sixth-order moment between PDFs. The results presented in Tab. I show that discrepancies are never larger than 3%, which in practice can be considered negligible. It is interesting to notice the remarkable agreement at least down to $p(z) = 10^{-6}$ (typical limit used in finance, see Ref. [4]).




FIG. 5. PDFs for a $q_n = 1$ noise and typical values of $(b; c)$ pair. a) (0.1; 0.1), $q = 1.021$ ($\chi^2 = 2.35 \times 10^{-9}$); b) (0.1; 0.88), $q = 1.287$ ($\chi^2 = 4.59 \times 10^{-10}$); c) (0.4; 0.1), $q = 1.26$ ($\chi^2 = 2.44 \times 10^{-9}$); d) (0.4; 0.4), $q = 1.38$ ($\chi^2 = 3.22 \times 10^{-7}$).




FIG. 6. PDFs for a $q_n = 1.2$ noise and typical values of $(b; c)$ pair. a) (0.1; 0.1), $q = 1.211$ ($\chi^2 = 8.67 \times 10^{-10}$); b) (0.1; 0.5), $q = 1.221$ ($\chi^2 = 6.01 \times 10^{-10}$); c) (0.3; 0.25), $q = 1.310$ ($\chi^2 = 8.11 \times 10^{-9}$); d) (0.3; 0.45), $q = 1.35$ ($\chi^2 = 7.36 \times 10^{-9}$).

B. Stationary distribution for (squared) volatility

The good results of the proposal presented in the previous sub-section have their explanation in the main feature of the ARCH processes, namely the temporal depen-

dence of the variance. This fact allow us to think of them along the same lines of Wilk and Włodarczyk [25] and of Beck [26]. They essentially considered a system following a Langevin equation with fluctuating temperature, genesis of the *superstatistics* (statistics of statistics) recently advanced by Beck and Cohen [15]. This approach was developed to treat non-equilibrium systems composed, for instance, of smaller cells obeying BG statistics with a distribution $P_{BG} \propto e^{-\beta E}$, E being the energy. The long-term stationary state that presents a spatio/temporal fluctuating temperature following a distribution $f(\beta)$. In the long-term, the probability density for the nonequilibrium system comes from BG statistics associated with the small cells that are averaged over the various β , i.e.,

$$P_{stationary}(E) = \int f(\beta) P_{BG}(E) d\beta. \quad (19)$$

Instead of defining the intensive fluctuating parameter, β , as the inverse temperature, we will define it as an inverse second-order moment, $\beta_\sigma = \frac{1}{2\sigma^2}$. For long times, we can study *GARCH* as a stationary diffusion process, composed by t increments, each of them of size z_t . The total value is given by $x_t \equiv \sum_{i=1}^t z_i$, where each increment follows a certain q_n -Gaussian, with a second-order moment, σ^2 , that is associated with a probability distribution, $p_\sigma(\sigma^2)$. The stationary probability distribution $p(z)$ is thus given by

$$p(z) = \int_0^\infty p_\sigma(\sigma^2) p(z|\sigma^2) d(\sigma^2), \quad (20)$$

where $p(z|\sigma^2)$ is the *conditional* probability of having a value z for the return given a value σ^2 . The homoskedastic case corresponds to $p_\sigma(\sigma^2) = \delta(\sigma^2 - \bar{\sigma}^2)$. Let us focus, for now, on the case $q_n = 1$ (Gaussian noise) with

$$p(z|\sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{z^2}{2\sigma^2}}. \quad (21)$$

In their proposal Beck and Cohen [15] showed that, if the intensive parameter β is associated with a Gamma distribution, then the macroscopic non-equilibrium steady state follows *exactly* a q -exponential distribution (see also [25,26]). Following the reverse line and assuming $p(z) = P_{stationary} \simeq P(z)$, we are lead to a Gamma distribution in β_σ . In other words,

$$p_\sigma(\sigma^2) = \frac{\exp\left(-\frac{1}{2\kappa\sigma^2}\right) (\sigma^2)^{-1-\lambda}}{(2\kappa)^\lambda \Gamma[\lambda]}, \quad (22)$$

where

$$\lambda = \frac{1}{q-1} - \frac{1}{2} \quad (23)$$

and

$$\kappa = \frac{1-q}{\bar{\sigma}^2(3q-5)}. \quad (24)$$

As can be seen in Fig. 7, the ansatz gives also a quite satisfactory description for the probability distribution in the (squared) volatility, corroborating the connection between the ARCH class of processes, nonextensive statistical mechanics, and superstatistics. The analytic expressions (22), (23) and (24) can be very useful in applications related, among others, to option prices [4,33] where volatility forecasting plays a particularly important role [34,35].




FIG. 7. The symbols in black represent the cumulative probability distribution, $P_{\geq}(\sigma^2)$ obtained from numerical simulation for a Gaussian noise with $b = c = 0.4$. The gray line represents the same distribution as given by Eq. (22) with $(\kappa, \lambda, \bar{\sigma}^2) = (0.444, 2.125, 1)$ satisfying Eqs. (23) and (24).

For the case of a q_n -Gaussian noise (with $q_n > 1$) similar arguments can be used. However, the achievement of an analytical solution for Eq. (20) which generalises the approach above is not a trivial task.

For the conditional probability $p(z|\sigma^2)$ we know a satisfactory answer [36], namely

$$p(z|\sigma^2) = \frac{\mathcal{A}_{(q_n, \sigma^2)}}{[1 + \mathcal{B}_{(q_n, \sigma^2)} (q_n - 1) z^2]^{\frac{1}{q_n - 1}}}. \quad (25)$$

But the same does not happen with $p_\sigma(\sigma^2)$. Assuming a continuous approach in q_n , a good ansatz for describing $p_\sigma(\sigma^2)$ is

$$\begin{aligned} p_\sigma(\sigma^2) &\propto (\sigma^2)^{-1-\lambda} \exp_{q_\sigma} \left(-\frac{1}{2\kappa\sigma^2} \right) \\ &= (\sigma^2)^{-1-\lambda} \left(1 + \frac{q_\sigma - 1}{2\kappa} \frac{1}{\sigma^2} \right)^{\frac{1}{1-q_\sigma}}, \end{aligned} \quad (26)$$

where q_σ is an index which depends on q_n , i.e., $q_\sigma(q_n)$ such that $q_\sigma(1) = 1$. For large σ^2 ,

$$p_\sigma(\sigma^2) \sim (\sigma^2)^{-1-\lambda}. \quad (27)$$

The integral $\int_0^\infty \sigma^2 p_\sigma(\sigma^2) d(\sigma^2)$ should equal the mean value $\bar{\sigma}^2 = \frac{a}{1-b-c}$. This yields,

$$\frac{1 + \lambda - \lambda q_\sigma}{2 \kappa (\lambda - 1)} = \bar{\sigma}^2, \quad (28)$$

and

$$\int_{-\infty}^{\infty} z^4 \int_0^{\infty} p_\sigma(\sigma^2) p(z | \sigma^2) d(\sigma^2) dz = 3 (\bar{\sigma}^2)^2 \frac{3q-5}{5q-7}. \quad (29)$$

From Eq. (27) and by adjusting numerically the curves for the cumulated probability distributions, $P_{\geq}(\sigma^2)$, we were able to determine λ , and then q_σ and κ from Eqs. (28) and (29). The procedure appears to be valid for values of q_n close to unity. From the analysis of some values of q_n we verified that the value of q_σ equals 1 for every q_n considered. Figures 7 and 8 confirm that our proposal produces a satisfactory approximation when compared with numerical simulations, particularly for large values of volatility (which are, in turn, responsible for the large returns). For the q_n -Gaussian noise case, although some discrepancy exists for small σ^2 , the tail is remarkably good.




FIG. 8. The line in black represent the cumulative probability distribution, $P_{\geq}(\sigma^2)$ obtained from numerical simulation for a q_n -Gaussian noise ($q_n = 1.15$) with $(b, c) = (0.5, 0)$. The gray line represents the same distribution as given by Eq. (26) with $(\kappa, \lambda, \bar{\sigma}^2) = (0.365, 2.371, 1)$, and $q_\sigma = 1$.

IV. DEGREE OF DEPENDENCE BETWEEN SUCCESSIVE RETURNS

As stated in Sec. II, stochastic variables, $\{z_t\}$, in a *GARCH* process are uncorrelated (see Eq. (1)). However, if we combine Eqs. (1) and (3), we can verify that they are not independent. More specifically, for *GARCH*(1, 1) ($\bar{\sigma}^2 = 1$), we have

$$z_t = \sqrt{1 + b(z_{t-1}^2 - 1) + c\left(\frac{z_{t-1}^2}{\omega_{t-1}^2} - 1\right)} \omega_t. \quad (30)$$

One of the possible measures of the dependence between the $(z_t; z_{t-1})$ stochastic variables consists in using a generalised form of the Kullback-Leibler relative entropy [37], namely

$$I_{q'}(p_1, p_2) \equiv - \int p_1(u) \ln_{q'} \left[\frac{p_2(u)}{p_1(u)} \right] du, \quad (31)$$

where $\ln_{q'}(x) \equiv \frac{x^{1-q'} - 1}{1-q'}$ (q' -logarithm). In the limit $q' \rightarrow 1$, $I_{q'}$ recovers the standard Kullback-Leibler form [38]. This generalised relative entropy equals zero whenever $p_2(u) = p_1(u)$, and has the same sign as q' otherwise. With these properties we can use $I_{q'}$ (with $q' > 0$) as a way to compute the “distance”, in probability space, from $p_2(u)$ to $p_1(u)$. Assume that u is a two dimensional random variable $u \equiv (x, y)$ so that $p_1(x, y)$ represents the *joint distribution* of (x, y) , and $p_2(x, y) \equiv h_1(x) h_2(y)$, where $h_1(x) \equiv \int p_1(x, y) dy$ and $h_2(y) \equiv \int p_1(x, y) dx$ are the *marginal distributions*. Random variables x and y are independent if $p_1(x, y) = p_2(x, y)$. With the functional $I_{q'}$ we can measure the degree of dependence *via* distance between probabilities $p_1(x, y)$ and $p_2(x, y)$.




FIG. 9. Dependence criterion $R_{q'}$ vs. q' for various *GARCH*(1,1) process with typical (q_n, b, c) triplets. In (a) $q_n = 1$ and (b, c) have been chosen as follows: 1-(0, 0), 2-(0.05, 0), 3-(0.1, 0.2), 4-(0.15, 0), 5-(0.2, 0), 6-(0.2, 0.2), 7-(0.25, 0), 8-(0.3, 0), 9-(0.4, 0), 10-(0.4, 0.1), 11-(0.4, 0.2), 12-(0.4, 0.4), 13-(0.5, 0), 14-(0.2, 0.688), 15-(0.35, 0), 16-(0.1, 0). In (b) $q_n = 1.2$ and (b, c) have been chosen as follows: 1-(0, 0), 2-(0.1, 0), 3-(0.1, 0.1), 4-(0.15, 0), 5-(0.2, 0), 6-(0.25, 0), 7-(0.3, 0.1), 8-(0.377, 0), 9-(0.3, 0.45), 10-(0.48, 0.0). The insets contain the derivative $dR_{q'}/dq'$ (numerically obtained) for the first four curves as mere illustration.

For this case, $u \equiv (x, y)$ and $q' > 0$, it was shown [40] that, besides a lower bound $I_{q'} = 0$ (total independence of x and y), $I_{q'}$ presents, for every value of q' , an upper bound (complete dependence between x and y) given by,

$$I_{q'}^{MAX}(p_1, p_2) = - \int \int [p_1(x, y)]^{q'} \{ \ln_{q'} h_1(x) + (1 - q') [\ln_{q'} h_1(x)] [\ln_{q'} h_2(y)] \} dx dy. \quad (32)$$

Dividing Eq. (31) by (32) we define

$$R_{q'} \equiv \frac{I_{q'}}{I_{q'}^{MAX}} \in [0, 1]. \quad (33)$$

This ratio can be used as a criterion for measuring the degree of dependence between random variables. Indeed, it presents an optimal q' , noted q^{op} [39], for which the sensitivity of $R_{q'}$ is maximal (q^{op} corresponds to the inflexion point of $R_{q'}(q')$). Higher (lower) values of q^{op} represents lower (higher) degree of dependence [40]. Taking $x = z_t$ and $y = z_{t-1}$ generated from Eq. (30), and applying Eq. (33), we obtained the curves presented in Fig. 9 for typical values of (b, c, q_n) . For each set we determined q^{op} and plotted it *versus* q obtained from Eq. (17): see Fig. 10. For both noises that have been illustrated, q^{op} monotonically decreases with q . For fixed q_n , this (decreasing) curve does not depend on the values of (b, c) . An illustration of this independence is indicated in Fig. 10 by using different pairs (b, c) that give the same q . This fact suggests the existence of a relation between non-Gaussianity (represented by q), the degree of dependence quantified with q^{op} , and the noise index q_n . This triangular relation (q, q^{op}, q_n) is analogous to another one which could relate the sensitivity, relaxation and stationarity in weakly chaotic systems such as those in which long-range interactions are assumed [41].




FIG. 10. Plot of q^{op} vs. q for typical (q_n, b, c) triplets. The arrow points two examples which were obtained from *different* triplets, and nevertheless coincide in what concerns the resulting point (q, q^{op}) .

V. CONCLUDING REMARKS

In this article, we presented a study of the stationary statistical features of the $GARCH(1, 1)$ stochastic pro-

cess, which is characterised by the fact that it exhibits a time dependent (and correlated) volatility. This attribute makes this system similar to those exhibiting fluctuations in some intensive parameter (e.g. temperature), which is at the basis of superstatistics. Inspired by the intimate connection between superstatistics and nonextensive statistical mechanics, we obtained an expression linking the dynamical parameters (b, c) and the noise nature, q_n , with the entropic index q characterising the stationary distribution for the associated $GARCH(1, 1)$ process. After numerically testing the validity of the approach for the return distributions, we derived analytical expressions for the squared volatility stationary distribution in the presence of q_n -Gaussian noise. The results are satisfactory for $q_n \simeq 1$. Then, using the q -generalised Kullback-Leibler relative entropy, we quantified the degree of dependence between successive returns. This analysis led to an entropic index q^{op} which is optimal in the sense that the ratio (33) exhibits maximal sensitivity. We then verified the existence of a direct relation between q^{op} , the non-Gaussianity, q , and the nature of the noise, q_n . An interesting property emerged, namely that, whatever be the pair (b, c) that results in a certain q for the stationary distribution, one obtains the same value of q^{op} . Consequently, the time series will present the same degree of dependence. The connection between various entropic indices reminds us of the dynamical scenario within which nonextensive statistical mechanics is formulated. Indeed, various entropic indices emerge therein, which coalesce onto the same value $q = 1$ if ergodicity is satisfied. In the present context, this connection can be analysed as follows. Due to the dynamical characteristics of $GARCH(1, 1)$, non-Gaussian distribution for returns ($q \neq 1$) comes from temporal dependence on its second-order moment which is a self-correlated variable. This correlation, sign of memory in the process, is the responsible for the breakdown of independence [19] between z_t and z_{t+1} , which in turn reflects on the entropic index q^{op} . The connection between q and q^{op} opens the door to the establishment of a relation between the topologies of phase space and probability space. A careful analysis of other kind of systems (e.g. long-range hamiltonian models, Langevin-like dynamics obeying generalised correlated Fokker-Plank equations) should give a deep insight onto this question.

L.G. Moyano is acknowledged for fruitful comments on the manuscript. Partial financial support from Faperj, CNPq, PRONEX/MCT (Brazilian agencies) and FCT/MES (Portuguese agency; contract SFRH/BD/6127/2001) is acknowledged as well.

[1] S. Abe and N. Suzuki, Europhys. Lett. **65**, 581 (2004).

- [2] M. Ausloos and K. Ivanova, Phys. Rev. E **63**, 047201 (2001).
- [3] O.A. Rosso, M.T. Martin, A. Plastino, Physica A **320**, 497 (2003).
- [4] R. Osorio, L. Borland and C. Tsallis, in *Nonextensive Entropy - Interdisciplinary Applications*, eds. M. Gell-Mann and C. Tsallis (Oxford University Press, New York, 2004).
- [5] S.M. Duarte Queirós, Physica A **344**, 279 (2004).
- [6] J.P. Bouchaud and M. Potters, *Theory of Financial Risks: From Statistical Physics to Risk Management* (Cambridge University Press, Cambridge, 2000); R.N. Mantegna and H.E. Stanley: *An introduction to Econophysics: Correlations and Complexity in Finance* (Cambridge University Press, Cambridge, 1999).
- [7] R.F. Engle, Econometrica **50**, 987 (2002).
- [8] F. Black and M. Scholes, J. Political Economy **81**, 637 (1973).
- [9] T. Bollerslev, R.Y. Chou and K.F. Kroner, J. Econometrics **52**, 5 (1992).
- [10] S.H. Poon and C. W. J. Granger, J. Econ. Lit. **41**, 478 (2003).
- [11] T. Bollerslev, J. Econometrics **31**, 307 (1986); S. J. Taylor, *Modelling Financial Time Series* (Wiley, New York, 1986).
- [12] F. Baldovin, Physica A, **342**, 119 (2004).
- [13] S.M. Duarte Queirós and C. Tsallis, Europhys. Lett. (2005), in press [cond-mat/0401181].
- [14] U. Frisch, *Turbulence: The Legacy of A. Kolmogorov*, (Cambridge University Press, Cambridge, 1997).
- [15] C. Beck and E. G. Cohen, Physica A **322**, 267 (2003).
- [16] P. Embrechts, C. Kluppelberg and T. Mikosch, *Modelling Extremal Events for Insurance and Finance (Applications of Mathematics)*, (Springer-Verlag, Berlin) 1997.
- [17] C. Tsallis, J. Stat. Phys. **52**, 479 (1988). A regularly an updated bibliography on the subject is available at <http://tsallis.cat.cbpf.br/biblio.htm>.
- [18] For systems presenting equiprobability whose (allowed) phase space, Ω , grows like $\Omega \sim N^\rho$ ($\rho > 0$), if we consider $q = 1 - \frac{1}{\rho}$, S_q will be proportional to N (and so *extensive*) in the same way S_{BG} is for systems with $\Omega \sim \mu^N$ ($\mu > 1$), respecting Clausius concept on macroscopic entropy and thermodynamics. In a probability context this can be related with an amendment in the law of probability of a system composed of N independent parts. A sheer analysis on this question can be found in Ref. [19,20].
- [19] C. Tsallis, Proceedings of the 31st Workshop of the International School of Solid State Physics “Complexity, Metastability and Nonextensivity”, held at the Ettore Majorana Foundation and Centre for Scientific Culture (Erice, July 2004), eds. C. Beck, A. Rapisarda and C. Tsallis (World Scientific, Singapore, 2005), in press [cond-mat/0409631].
- [20] Y. Sato and C. Tsallis, Proceedings of the Summer School and Conference on Complexity (Patras and Olympia, July 2004), ed. T. Bountis, International Journal of Bifurcation and Chaos (2005), in press [cond-mat/0411073].
- [21] C. Tsallis, M. Gell-Mann and Y. Sato, *Special scale-invariant occupancy of phase space makes the entropy S_q additive*, preprint (2005).
- [22] M. L. Lyra and C. Tsallis, Phys. Rev. Lett. **80**, 53 (1998); F. Baldovin and A. Robledo, Europhys. Lett. **60**, 518 (2000).
- [23] L. Borland, Phys. Rev. E **57**, 6634 (1998).
- [24] C. Beck, G.S. Lewis and H.L. Swinney, Phys. Rev. E **63**, R035303 (2001).
- [25] G. Wilk and Z. Włodarczyk, Phys. Rev. Lett. **84**, 2770 (2000).
- [26] C. Beck, Phys. Rev. Lett **87**, 180601 (2001).
- [27] V. Latora, A. Rapisarda and C. Tsallis, Phys. Rev. E **64**, 056134 (2001).
- [28] M. Gell-Mann and C. Tsallis, eds., *Nonextensive Entropy - Interdisciplinary Applications* (Oxford University Press, New York, 2004).
- [29] C. Tsallis, A.R. Plastino and R.S. Mendes, Physica A **261**, 534 (1998).
- [30] D. Prato and C. Tsallis, Phys. Rev. E **60**, 2398 (1999).
- [31] C. Tsallis, Braz. J. Phys **29**, 1 (1999).
- [32] A.M.C. de Souza and C. Tsallis, Physica A **236**, 52 (1997).
- [33] L. Borland, Phys. Rev. Lett. **89**, 098701 (2002); L. Borland, Quantitative Finance **2**, 415 (2002).
- [34] P. Willmott, *On Quantitative Finance*, 2nd edition (John Wiley & Sons, New York, 2000).
- [35] L. Borland and J.P. Bouchaud, [cond-mat/0403022].
- [36] Since σ_t and ω_t are uncorrelated and independent, for a given value σ , we can do a change of variable, Eq. (1), which transforms Eq. (16) into the conditional probability Eq. (25).
- [37] C. Tsallis, Phys. Rev. E **58**, 1442 (1998).
- [38] S. Kullback and R.A. Leibler, Ann. Math. Stat. **22**, 79 (1961).
- [39] Optimal in the sense that its two extreme values correspond to full independence/dependence [40].
- [40] L. Borland, A.R. Plastino and C. Tsallis, J. Math. Phys. **39**, 6490 (1998) [Erratum: J. Math. Phys. **40**, 2196 (1999)].
- [41] C. Tsallis, Physica A **340**, 1 (2004).

q_n	b	c	$\langle z^6 \rangle_{\text{numerical}}$	$\langle z^6 \rangle_{\text{ansatz}}$	error (%)
1	0.1	0.2	13.97	14.05	0.60
1	0.1	0.88	66.75	65.46	1.93
1	0.4	0.1	61.45	62.33	1.44
1	0.4	0.4	591.91	517.67	1.75
1.2	0.1	0.1	49.41	49.08	0.67
1.2	0.1	0.5	55.93	55.43	0.89
1.2	0.3	0.25	181.48	179.44	1.13
1.2	0.3	0.45	1416.41	1455.37	2.75

TABLE I. Percentual error in the *sixth*-order moment between numerical and ansatz PDFs presented in Figs. 5 and 6.